

SOXS




Son Of X-Shooter at NTT

Sergio Campana

Osservatorio astronomico di Brera

On behalf of a large collaboration

What is X-shooter



UVB			VIS			NIR		
Slit width (")	Resolution ($\lambda/\delta\lambda$)	Sampling (pix/FWHM)	Slit width (")	Resolution ($\lambda/\delta\lambda$)	Sampling (pix/FWHM)	Slit width (")	Resolution ($\lambda/\delta\lambda$)	Sampling (pix/FWHM)
0.5	9100	3.5	0.4	17400	3.0	0.4	11300	2.0
0.8	6300	5.2	0.7	11000	4.8	0.6	8100	2.8
1.0	5100	6.3	0.9	8800	6.0	0.9	5600	4.0
1.3	4000	8.1	1.2	6700	7.9	1.2	4300	5.3
1.6	3300	9.9	1.5	5400	9.7	1.5	3500	6.6
IFU	7900	4.1	IFU	12600	4.2	IFU	8100	2.8

Band	U	B	V	R	I	J	H	K'
mag	21.5	21.7	21.7	21.6	21.2	20.5	20.8	19.3

Continuum spectrum S/N=10 - 1 hr exposure

NOT Transient Explorer – A new work-horse for the Nordic Optical Telescope



A cross-dispersed spectrograph covering 350-1700 nm, resolution ~ 4000 (possibly with also a higher-res mode), single slit (with different choices for the slit width), including ADC and efficient enough to be sky-limited in 30 min integration.



Visible imager with 5-6 arcmin FOV, 2k x 2k detector, sampling 0.15-0.18 arcsec per pixel.



Near-IR imager using a 2k x 2k HAWAII-II detector with same FOV and sampling as in the visible.





De-scoped version: imaging reduced to a visible slit-viewing camera with FOV of 3 arcmin (similar to StanCam).

Nordic (Denmark Sweden, etc.) + Italian collaboration




SOXS Science case: the transient sky



Just a few science cases


- Minor planets and asteroids
- Young stellar objects
- Planetary transits
- X-ray binary transients
- Novae
- Supernovae (Ia, CC)
- GRB
- GW-&neutrino EM counterparts
- Radio sky transients & fast radio bursts

Discoveries




Water vapor in the atmosphere of a transiting planet

First SN shock break out



Major outburst 2 yr before the (probable) SN explosion



The most distant object in the Universe (at the time of discovery)

A working example

During 2005-2013 Nature published ~ 180 astronomical papers with more than 50 citations.

Among them **36%** are on transients objects.

PESSTO

An already working example

- ~20% of selected candidates from SN searches enter into the observing queue
- ~ 50% of the transients are eventually observed and classified


90% remain unclassified

GAIA Transient Alerts

GAIA
is
coming

	<i>alerting object</i>	<i>5-yrs (Entire Mission)</i>	<i>main location</i>
interesting	Supernovae <19 mag	6000	out of plane
	Microlensing (bulge)	~1000	bulge/plane
	Microlensing (all sky)	~700	out of plane
	GRB optical counterparts	~hundreds (?)	out of plane
	R CrB-type stars	~hundreds (?)	gal. plane
	CN	150	gal. plane
	FU Ori	14	gal. plane
contaminants(?)	Eclipsing binaries	a million (?)	gal. plane
	AGNs	500,000 (?)	out of plane
	Asteroids	thousands (?)	out of plane
	Be stars	thousands (?)	gal. plane
	Long period variables/Miras	thousands (?)	gal. plane
	M-dwarf flares	2000	gal. plane
	DN (U Gem) (except rare big flares)	500 (?)	gal. plane

What is SOXS



Spectroscopic machine for the transient sky.


Even now with PESSTO in place $>70\%$ of newly discovered transients remain without spectroscopic follow-up.

In the near future years there will be many imaging survey wide-field telescopes (iPTF, DES, Pan-STARRS, LSST) as well as high-energy transients (Swift, INTEGRAL, MAXI), GAIA-alters GW-alters etc. but very limited spectroscopic follow-up


SOXS @ NTT

We propose to build and operate a spectroscopic facility, SOXS (Son of X-Shooter), with a wide spectral coverage (0.35-1.75 μm) and good spectral resolution ($R \sim 4,500$) able to characterize and follow-up in depth any kind of transient source


A possible optical layout of the Common Path



SOXS @ NTT



Initial performances




BLUE arm


0.5 arcsec box

0 and ± 12 arcsec
positions


BLUE spectrograph



Pixel size 15 micron
2048x2048



Initial performances




RED arm


0.5 arcsec box

0 and ± 12 arcsec
positions

RED spectrograph



Pixel size 18 micron
2048x2048



SOXS performances


- Still TDB
- Goal:

continuum spectrum $R \sim 20$ -20.5 $S/N=10$ in 1 hr

This nicely match limiting magnitude of current (e.g. iPTF) synoptic surveys

“Extended” guiding camera to use as imaging instrument >3 arcmin FOV.

Consortium structure



Science Board

S. Campana (INAF-OABrera) - Italy

E. Cappellaro (INAF-OAPadova) - Italy

M. Della Valle (INAF-OANapoli) - Italy

A. De Ugarte Postigo (IAA-CSIS) - Spain

J. Fynbo (Dark-NBI) - Denmark

M. Hamuy (Millenium Inst.) - Chile

G. Pignata (Millenium Inst.) - Chile

S. Smartt (Univ. Belfast) – UK

...

Timeline 2016-2020

Project Phase	Start	End	Duration
Phase A	05/2015	09/2015	5 months
Phase B	10/2015	02/2016	5 months
Phase C	03/2016	12/2016	10 months
Phase D	01/2017	06/2019	30 months
Phase E	07/2019	>06/2022	TBD

but...

**ESO decision for phase A approval
in April 2015**

Operations

ESO will reward the consortium with NTT observing time in the 2016-2020 time-frame.

We will start from 2016 with existing instruments (EFOOSC2+SOFI) and when SOXS will be ready (mid-2019) we will continue with SOXS.

We will likely have ~ 150 n/yr.

Prospects to continue beyond 2020 (GW, LSST, etc.)

Source class	Obs. Time	Key project & Aim
All	500 <u>hr</u>	Fast characterization of transients from other surveys
Open	500 <u>hr</u>	Open time for spectroscopic <u>ToO</u> observations
Asteroids & TNO	200 <u>hr</u>	Characterization of populations of minor bodies, input to models of <u>solar</u> system formation and mitigation of impact hazard
Comets and new comets	100 <u>hr</u>	Monitor of >5 bright stars for primary and secondary eclipses
Planetary transits	200 <u>hr</u>	
Young stellar objects	100 <u>hr</u>	
Stars	100 <u>hr</u>	
X-ray binary transients	200 <u>hr</u>	Derive the mass function of >10 <u>XRB</u> transients in outburst
<u>Magnetars</u>	50 <u>hr</u>	Fast follow up of >10 <u>magnetar's</u> flares
Novae	100 <u>hr</u>	Statistical sample of >150 <u>SNe Ia</u> in the low- z Universe to study the <u>local</u> properties and dust extinction
ILOT	300 <u>hr</u>	
SN <u>Ia</u>	500 <u>hr</u>	
CC-SN	500 <u>hr</u>	
Super-luminous supernovae	500 <u>hr</u>	Build a statistical spectroscopic sample of <u>SLSN</u>
Prompt GRB	100 <u>hr</u>	Fast spectroscopy of >50 GRBs to probe the galaxy host medium
High- z ($z>5$) GRB	50 <u>hr</u>	Transmission spectra of >5 high-redshift GRBs
<u>GRB-SNe</u>	100 <u>hr</u>	Follow the evolution of >5 SN associated to nearby ($z<0.3$) GRBs
Active galactic nuclei and <u>blazars</u>	200 <u>hr</u>	Study the spectral evolution of >10 TDEs
Tidal disruption events	100 <u>hr</u>	
Gravitational Wave triggers	200 <u>hr</u>	Spectroscopic follow up of candidate GW counterparts. This includes <u>kilonovae</u> from short GRBs.
Neutrino triggers	100 <u>hr</u>	Spectroscopic follow up of candidate neutrino counterparts
Unknown	300 hr	

Data policy

10% of the consortium time open to the community as fast ToO (Swift-like) observations (public data)

Relevant information (redshift, peculiar sources, etc.) announced in real time through GCN, ATEL, IAUC, etc.

Consortium data TDB if public from the beginning or with some short (1 months) proprietary period.

Funds

>60% secure funds (as most of the projects)

Remaining funds to be asked for if proposal approved

If you are interested (and have funds and/or hardware capabilities) we are open to collaboration

(deadline February 13)

**If you want to support the project send
me an e-mail**

**[http://www.brera.inaf.it/~campana/
SOXS/Supporting.html](http://www.brera.inaf.it/~campana/SOXS/Supporting.html)**

The more we are, the best it is!

Thanks