




♦ INAF  
ISTITUTO NAZIONALE  
DI ASTROFISICA  
NATIONAL INSTITUTE  
FOR ASTROPHYSICS



Smithsonian  
Institution



# Paolo Esposito

The background of the slide features a deep space scene with a large, bright blue nebula on the right and a smaller, smaller spiral galaxy with a bright core and a distinct spiral arm on the left.

## The CATS @ BAR WR/BH X-ray binaries

+ **G. L. Israel**, L. Sidoli, L. Zampieri,  
M. Mapelli, D. Milisavljevic, G. Fabbiano

CNOC IX  
Sep 23, 2015

# Thousands of X-ray unidentified sources

CXO DFS - CXC/NASA/Giacconi



Faint X-ray sources can remain  
unidentified for years


~50,000 objects from past missions

>500,000 with XMM, Chandra, Swift

- Different populations of X-ray sources
- Interesting srcts might lurk among them





Periodic signals  
are key to  
understand the  
nature of a  
source!



# X-ray pulsators

In general, modulation is discovered through timing analysis of the source targeted by an observation



Open stellar cluster: Praesep

European Space Agency


**Enormous discovery space**  
in serendipitous sources

# Highlights from GLI's previous searches

- **EXOSAT:** 4U 0142+614, prototype of the AXP class (I+94)
- ROSAT PSPC: HD 49798, a very massive WD in a post common envelope phase (I+96, Mereghetti+09)
- ROSAT HRI: the 2-WD system HM Cnc (I+99,02; E+14), the binary with the shortest orbital period known: 5.4 min!

---



AXPs and SGRs host **magnetars**, NSs with  $B \sim 10^{15}$  G



Ramifications in all directions and across many fields

# Highlights from GLI's previous searches

- EXOSAT: 4U 0142+614, prototype of the AXP class (I+94)
- **ROSAT PSPC: HD 49798, a very massive WD in a post common envelope phase** (I+96, Mereghetti+09)
- ROSAT HRI: the 2-WD system HM Cnc (I+99,02; E+14), the binary with the shortest orbital period known: 5.4 min!



A promising candidate precursor of SN Ia


# Highlights from GLI's previous searches

- EXOSAT: 4U 0142+614, prototype of the AXP class (I+94)
- ROSAT PSPC: HD 49798, a very massive WD in a post common envelope phase (I+96, Mereghetti+09)
- **ROSAT HRI**: the **2-WD system HM Cnc** (I+99,02; E+14), **the binary with the shortest orbital period known: 5.4 min!**

---

HM Cnc: the **brightest persistent Galactic source of GWs!**

GSFC/D. Berry




RX J0806.3+1527  
aka HM Cnc



aka 'The Stinker'


SDO/GLI




A new emission mechanism (**UIM**) possibly identified

# Individual sources


Normalized Count Rate



# Individual sources




# CXOU J123030.3+413853 in NGC 4490





- Multiple CXO observations
- Modulation at  $P = (6.4 \pm 0.1)$  h  
(confirmed also by XMM)
- Pulsed fraction: ~90 per cent

(First observed by Roberts et al. 2002)



# CXOU J123030.3+413853 in NGC 4490



- Multiple CXO observations
- Modulation at  $P = (6.4 \pm 0.1)$  h  
(confirmed also by XMM)
- Pulsed fraction: ~90 per cent

(First observed by Roberts et al. 2002)

# A 6.4 hr BH binary in NGC 4490


NGC 4490 is a spiral galaxy  
interacting with the irregular  
NGC 4485

$P = 6.4$  h, 90% PF




~2.5 kpc from NGC 4490's  
nucleus

$L$  from ~0.2 to  $2 \times 10^{39}$  erg s<sup>-1</sup>

$L > L_E$  for a  $5\text{--}10 M_\odot$  object  
(similar lower limit from diskbb fit)



# Circinus Galaxy X-1



Bauer et al. 2001:  $P \sim 7.2 \text{ h}$

$$L_X = (1-5) \times 10^{40} \text{ erg s}^{-1}$$

See Weisskopf+2004


Strictly speaking, not a CATS@BAR source, we sort of bumped into it

D = 4.2 Mpc

# NGC 4490's J123030 & CG X-1

$P = 6.4 / 7.2$  h, ~90% PF

$L \sim 10^{38/39}$  and  $10^{39/40}$  erg s<sup>-1</sup>




# NGC 4490's J123030 & CG X-1

P = 6.4 / 7.2 h, ~90% PF

L  $\sim 10^{38/39}$  and  $10^{39/40}$  erg s $^{-1}$

Foreground polars?



# NGC 4490's J123030 & CG X-1

$P = 6.4 / 7.2 \text{ h}$ ,  $\sim 90\%$  PF

$L \sim 10^{38/39}$  and  $10^{39/40} \text{ erg s}^{-1}$

Foreground polars?


**P < 0.03%**

no counterparts,  $nH >$  Gal.  $nH$

LMXBs:

- Transients;
- Very different pulse profiles!

low-amplitude orbital modulation (if any):  
sharp eclipse ingresses/egresses (small  
X-ray emitting regions), dips



# NGC 4490's J123030 & CG X-1

$P = 6.4 / 7.2 \text{ h}$ ,  $\sim 90\%$  PF

$L \sim 10^{38/39}$  and  $10^{39/40} \text{ erg s}^{-1}$

~~Foreground polars?~~

$P < 0.03\%$


no counterparts,  $nH >$  Gal.  $nH$

LMXBs:

- Transients;
- Very different pulse profiles!

low-amplitude orbital modulation (if any):  
sharp eclipse ingresses/egresses (small  
X-ray emitting regions), dips

**UNLIKELY!**



# NGC 4490's J123030 & CG X-1

$P = 6.4 / 7.2 \text{ h}$ ,  $\sim 90\%$  PF

$L \sim 10^{38/39}$  and  $10^{39/40} \text{ erg s}^{-1}$

Foreground polars?

$P < 0.03\%$


no counterparts,  $nH >$  Gal.  $nH$

LMXBs:

- Transients;
- Very different pulse profiles!

low-amplitude orbital modulation (if any):  
sharp eclipse ingresses/egresses (small  
X-ray emitting regions), dips

UNLIKELY!



# NGC 4490's J123030 & CG X-1

$P = 6.4 / 7.2 \text{ h}$ ,  $\sim 90\%$  PF

$L \sim 10^{38/39}$  and  $10^{39/40} \text{ erg s}^{-1}$

Foreground polars?

$P < 0.03\%$


no counterparts,  $nH >$  Gal.  $nH$

LMXBs:

- Transients;
- Very different pulse profiles!

low-amplitude orbital modulation (if any):  
sharp eclipse ingresses/egresses (small  
X-ray emitting regions), dips

UNLIKELY!



# NGC 4490's J123030 & CG X-1

$P = 6.4 / 7.2 \text{ h}$ ,  $\sim 90\%$  PF

$L \sim 10^{38/39}$  and  $10^{39/40} \text{ erg s}^{-1}$

Foreground polars?


$P < 0.03\%$

no counterparts,  $nH >$  Gal.  $nH$

- Transients;
- Very different pulse profiles!

low-amplitude orbital modulation (if any):  
sharp eclipse ingresses/egresses (small  
X-ray emitting regions), dips

UNLIKELY!



HMXBs:  
Possible only with a WR star

# Wolf–Rayet stars



HST/Judy Smith

- Final phase (prior to core collapse) of massive stars
- H envelope stripped away via stellar wind or close binary evolution, revealing products of CNO (WN) or He burning (WC)
- Strong emission lines, intense ( $10^{-5} M_{\odot} \text{yr}^{-1}$ ) and fast ( $\sim(1-3)\times 10^3 \text{ km s}^{-1}$ ) winds
- Compact,  $R < 2R_{\odot}$  for  $M \sim 20M_{\odot}$

# NGC 4490's J123030 & CG X-1

$P = 6.4 / 7.2 \text{ h}$ ,  $\sim 90\%$  PF

$L \sim 10^{38/39}$  and  $10^{39/40} \text{ erg s}^{-1}$

Foreground polars?

$P < 0.03\%$


no counterparts,  $nH >$  Gal.  $nH$

LMXBs:

- Transients;
- Very different pulse profiles!

low-amplitude orbital modulation (if any):  
sharp eclipse ingresses/egresses (small  
X-ray emitting regions), dips

UNLIKELY!




HMXBs:  
Possible only with a WR star

WR + BH


A NS cannot be excluded,  $L_x \sim 10^{39}$   
maybe  $10^{40} \text{ erg s}^{-1}$  for the NuSTAR  
one in M82 (Bachetti+14)

# Pulse profile




Carbon copy of the WR/HMXB  
light curves!

Cyg X-3  
Hjalmarsdotter+08




IC10 X-1  
Prestwich+07

# Pulse profile




Carbon copy of the WR/HMXB  
light curves!




NGC300 X-1  
Carpano+07

NGC 253  
Maccarone+14

# Pulse profile



Carbon copy of the WR/HMXB  
light curves!




Signature pulse profile produced by absorption and Compton scattering of the X-rays through the WR stellar wind (even for  $i \sim 30^\circ$ ; Zdziarski et al. 2010)

# Counterparts

$M_V$  from  $-2.5$  to  $-7$  for WRs

NGC 4490's J123030



3 rather bright  
and blue objects

A  $M_V = -6.4$   
 $B-V = -0.4$


B  $M_V = -4.8$   
 $B-V = -0.1$

C  $M_V = -3.0$   
 $B-V = -1.2$

CG X-1

HST:  $M_V > -6.8$

# Luminosity



A BH needs a disc to shine

$$M_{\text{BH}} \gtrsim 1.5 v_{w, 1000}^4 \delta^2 M_{\odot}$$

Carpano+2007; Illarionov & Sunyaev 1975

$$L_{\text{X}} \approx \eta \frac{\dot{M}_{\text{w}} c^2 G^2 M_{\text{BH}}^2}{a^2 v_{\text{w}}^4}$$

$\dot{M}_{\text{w}} = 10^{-5} M_{\odot} \text{ yr}^{-1}$  (e.g. Crowther 2007)  
 $v_{\text{w}} = 1000 \text{ km s}^{-1}$

$$L_{\text{X}} > 3 \times 10^{39} \text{ erg s}^{-1} \text{ for } M > 10 M_{\odot}$$

For CG X-1,  $L_{\text{X}}$  up to  $2 \times 10^{40} \text{ erg s}^{-1}$

If  $L_E$  holds,  $M > 75 M_{\odot}$  for a He or C/O donor

For  $75 M_{\odot}$ :  $L_{\text{X}} = 2 \times 10^{40} \text{ erg s}^{-1}$

Easily more with RLO

Canonical stellar-mass BH if moderately beamed / super-Edd

# WR/BH binaries

## Cyg X–3

$2 M_{\odot}$  NS or a  $3\text{--}5 M_{\odot}$  BH  
 $P = 4.8$  h,  $L \sim 10^{38}$  erg s $^{-1}$

## M101 ULX–1

$>5 M_{\odot}$  BH;  $P = 8.2$  d  
(Liu et al. 2013 )

+ NGC 4490 and CG X–1  
+ candidate in NCG 253:  
 $P \sim 15$  h,  $L \sim 10^{38}$  erg s $^{-1}$   
(Maccarone et al. 2014)


WR/BH binaries are the  
progenitors of 2BHs with  
merger time scales shorter  
than a Hubble time

## NGC300 X–1

$20 M_{\odot}$  BH;  $P = 32.3$  h  
(Carpano et al. 2007)

## IC 10 X–1

$33 M_{\odot}$  BH;  $P = 34.9$  h  
(Prestwich et al. 2007)



# WR/BH binaries

Theoretical predictions from population synthesis:  
 $0.4 < 20 < 1000$  events/yr for aLIGO or Virgo  
(Abadie+2010)

Host galaxy	Source	Period (h)	BH mass ( $M_{\odot}$ )	WR mass ( $M_{\odot}$ )	SFR ( $M_{\odot} \text{ yr}^{-1}$ )	$Z$ ( $Z_{\odot}$ )	$t_{\text{GW}}$ (Gyr)
IC 10	X-1	34.9	33	35	0.07	0.22	1.4
NGC 300	X-1	32.8	20	26	0.14	0.19	1.7
NGC 4490	CXO J123030 *	6.4	—	—	4.5	0.23	0.038
NGC 253	CXO J004732 *	14.5	—	—	4.0	0.24	0.33
Circinus	CG X-1 *	7.2	—	—	1.5	0.10	0.052
M 101	ULX-1	196.8	20	19	3.1	0.17	200
Milky Way	Cyg X-3	4.8	3	7	0.25	0.31	0.051

Natal kick? Mass of the 2<sup>nd</sup> compact object?

- WR forms a  $10 M_{\odot}$  BH via direct collapse
  - Orbital parameters left **unchanged**
  - If BH mass unknown, we assume  $10 M_{\odot}$

# WR/BH binaries

Host galaxy	Source	Period (h)	BH mass (M <sub>⊙</sub> )	WR mass (M <sub>⊙</sub> )	SFR (M <sub>⊙</sub> yr <sup>-1</sup> )	Z (Z <sub>⊙</sub> )	t <sub>GW</sub> (Gyr)
IC 10	X-1	34.9	33	35	0.07	0.22	1.4
NGC 300	X-1	32.8	20	26	0.14	0.19	1.7
NGC 4490	CXO J123030 *	6.4	—	—	4.5	0.23	0.038
NGC 253	CXO J004732 *	14.5	—	—	4.0	0.24	0.33
Circinus	CG X-1 *	7.2	—	—	1.5	0.10	0.052
M 101	ULX-1	196.8	20	19	3.1	0.17	200
Milky Way	Cyg X-3	4.8	3	7	0.25	0.31	0.051

$$R = \rho_{\text{SFR}}(z) \sum_i (t_{\text{GW}, i} + t_{\text{evol}, i})^{-1} (\text{SFR}_i)^{-1} \text{ yr}^{-1} \text{ Mpc}^{-3}$$

# WR/BH binaries

Host galaxy	Source	Period (h)	BH mass (M <sub>☉</sub> )	WR mass (M <sub>☉</sub> )	SFR (M <sub>☉</sub> yr <sup>-1</sup> )	Z (Z <sub>☉</sub> )	t <sub>GW</sub> (Gyr)
IC 10	X-1	34.9	33	35	0.07	0.22	1.4
NGC 300	X-1	32.8	20	26	0.14	0.19	1.7
NGC 4490	CXO J123030 *	6.4	—	—	4.5	0.23	0.038
NGC 253	CXO J004732 *	14.5	—	—	4.0	0.24	0.33
Circinus	CG X-1 *	7.2	—	—	1.5	0.10	0.052
M 101	ULX-1	196.8	20	19	3.1	0.17	200
Milky Way	Cyg X-3	4.8	3	7	0.25	0.31	0.051

$$R = \rho_{\text{SFR}}(z) \sum_i (t_{\text{GW}, i} + t_{\text{evol}, i})^{-1} (\text{SFR}_i)^{-1} \text{ yr}^{-1} \text{ Mpc}^{-3}$$

$$t_{\text{GW}} = \frac{5}{256} \frac{c^5 a^4 (1 - e^2)^{7/2}}{G^3 m_1 m_2 (m_1 + m_2)} \cdot \begin{aligned} & \rho_{\text{SFR}}(z) \text{ cosmic SFR density} \\ & t_{\text{evol}} \sim 3 \text{ Myr} \end{aligned}$$

(Peters 1964) (Mapelli+2010)

$$\rho_{\text{SFR}}(z = 0.3) \sim 3.6 \times 10^{-2} \text{ M}_\odot \text{ yr}^{-1} \text{ Mpc}^{-3} \quad (\text{Hopkins \& Beacom 2006})$$

# WR/BH binaries

Host galaxy	Source	Period (h)	BH mass (M <sub>⊙</sub> )	WR mass (M <sub>⊙</sub> )	SFR (M <sub>⊙</sub> yr <sup>-1</sup> )	Z (Z <sub>⊙</sub> )	t <sub>GW</sub> (Gyr)
IC 10	X-1	34.9	33	35	0.07	0.22	1.4
NGC 300	X-1	32.8	20	26	0.14	0.19	1.7
NGC 4490	CXO J123030 *	6.4	—	—	4.5	0.23	0.038
NGC 253	CXO J004732 *	14.5	—	—	4.0	0.24	0.33
Circinus	CG X-1 *	7.2	—	—	1.5	0.10	0.052
M 101	ULX-1	196.8	20	19	3.1	0.17	200
Milky Way	Cyg X-3	4.8	3	7	0.25	0.31	0.051

$$R = \rho_{\text{SFR}}(z) \sum_i (t_{\text{GW}, i} + t_{\text{evol}, i})^{-1} (\text{SFR}_i)^{-1} \text{ yr}^{-1} \text{ Mpc}^{-3}$$

$$t_{\text{GW}} = \frac{5}{256} \frac{c^5 a^4 (1 - e^2)^{7/2}}{G^3 m_1 m_2 (m_1 + m_2)} \cdot \begin{aligned} & \rho_{\text{SFR}}(z) \text{ cosmic SFR density} \\ & \cdot t_{\text{evol}} \sim 3 \text{ Myr} \end{aligned}$$

(Peters 1964) (Mapelli+2010)

**<16 yr<sup>-1</sup> for distance range 1 Gpc**

**0.4 <20 <1000 yr<sup>-1</sup> (Abadie+2010) / ~10 yr<sup>-1</sup> (1.4 Gpc) (Maccarone+2014)**

# The Future - I

## CATS @ BAR

- Every 3/4 months we run the pipeline on the new ACIS data; **new pulsators are coming**
- **Z<sup>2</sup>-like algorithms**, better suited for low-counts sources
  - X-ray/optical **follow-ups & classification**
  - Search the **XMM EPIC** archive (**6–73 ms sampling**) in the **EXTras** project



## NGC 4490's 6.4 h BH binary

- Identification of the **companion star**
- Dynamical measurement of the **BH mass**
- New estimates of the **metallicity** of NGC 4490/85

# The Future - II

- Dearth of WR HMXBs. Yet they should be relatively common and very bright. Are there others known systems which may have been misclassified?
- The sample is still small, but growing (from 3 to 7 in 2 yr!)
- Can a significant fraction of ULXs be powered by WRs?
- Stellar massive BHs  $>20 M_{\odot}$
- Population synthesis and common envelope phase
- Rate of (stellar) 2-BH mergers to be seen (Advanced Ligo)

